



Dallas Moore, MS, RPh 7 January 2021

©UNIVERSITY OF UTAH HEALTH

DATA – SENIORS AND MEDICATIONS

- From Health and Human Services:
 - More than 10,000 people turn 65 every day in the United States
 - Fifty-five percent of elderly are non-compliant with medication instructions
 - Approximately 200,000 older adults are hospitalized annually due to adverse drug reactions²

^{2: &}lt;u>Patient Safety Organizations: Hospital Participation, Value, and Challenges Report (OEI-01-17-00420) 09-25-2019 (hhs.gov)</u>



^{1:} Reducing Medication Problems of the Elderly: A Management Advisory Report (OEI-04-89-89122; 10/90) (hhs.gov)


DATA – SENIORS AND MEDICATIONS

- Lown Institute report
 - More than 40% older Americans regularly take five or more medications and almost 20% take more than 10 medications
 - Every day, approximately 750 older Americans are hospitalized due to serious side effects from one or more medications
 - the odds of experiencing a serious adverse reaction to a medication increases 7–10% with each additional drug
 - 1. Solving Medication Overload: America's Other Drug Problem | Health Affairs



MEDICATION OBSTACLES FOR SENIORS

- Common situations that may inhibit a senior from managing medications appropriately
- 1. Financial
- 2. Access to Pharmacy/Medications
- 3. Vision Loss
- 4. Memory Loss
- 5. Decreased Ability to Swallow
- 6. Hearing Loss
- 7. Social Isolation





FINANCIAL

 Problem: Can range from limited income to extremely expensive medications

- Use generics when possible
- Research finance assistance programs
 - Samples; Manufacture coupons; GoodRx;
- Purchase additional Medicare Part D plan



ACCESS TO PHARMACY/MEDICATIONS

 Problems: When multiple prescriptions, can't keep track; transportation to pharmacy

- Mailorder pharmacies
- Bubble/compliance packaging
- Convenience timing
 - Work with thirds party insurance to move all refill dates to once per month for all prescriptions



ACCESS TO PHARMACY/MEDICATIONS

 Problems: When multiple prescriptions, can't keep track; transportation to pharmacy

- Deprescribing
 - <u>Deprescribing.org</u> (Bruyere Research Institute Ontario, Canada)
 - Medication Appropriateness Index
 - Beers Criteria/List



VISION LOSS

 Problem: Not able to read small print on vials or distinguish between medications

- Ask if pharmacy can printer large font labels
- Bubble/convenience packaging
- Microchipped label or bottle with reader
 - Can speak the label information



MEMORY LOSS

 Problem: Can range from simple forgetfulness to Alzheimer's/Dementia

- Use pill boxes, bubble/convenience packaging, auto-dispenser
- Alarmed pill boxes
- Professional in-home caregivers



DECREASED ABILITY TO SWALLOW

Problem: Many seniors develop a problem of swallowing

- Identify medication to be crushed or mixed
- Convert to liquid



HEARING LOSS

 Problem: Patient may not have heard or understood administration instructions

- Utilize caregiver (family or professional)
- Share printed instructions
- Pharmacy to follow-up with patient to verify patient knows how to take correctly



SOCIAL ISOLATION

 Problem: Many elderly live alone with no support and may become disinterested or depressed

- Utilize family or professional caregiver
- Be aware

QUESTIONS/DISCUSSION



